Environmental Informatics Archives, Volume 3 (2005), 111 – 117
EIA05-015
ISEIS Publication Series Number P002
© 2005 ISEIS - International Society for Environmental Information Sciences

Photodegradation of Acid Blue 29 and Ethyl Violet in the Presence/Absence of Sodium Hydroxide and Aluminium Ions

C-C. CHEN[1*] A.J. Chaudhary[2] and S.M. Grimes[2]
[1]Center for Environmental, Safety and Health Technology Development, Industrial Technology Research Institute, W200, 195, Sec. 4, Chung Hsing Road, Chutung, Hsinchu, Taiwan 310, China
[2]Institute for the Environment, Brunel University, Uxbridge, London UB8 3PH, UK

Abstract. The photodegradation of Acid Blue 29 and Ethyl Violet in the presence/absence of sodium hydroxide (NaOH) and aluminium ions was investigated. The results show that the dyes degrade efficiently in the presence of 2 M NaOH. The addition of titanium dioxide as a catalyst and hydrogen peroxide as an oxidant, however, does not enhance the degradation rate of the dyes. The presence of aluminium ions (Al^{3+}) in the dye solution retards the degradation of the dyes.

Keywords: Photodegradation, Acid Blue 29, Ethyl Violet, NaOH, aluminium

1. Introdution

Dyes are used to colour products, mainly in the textile industry. Many dyes are inert and non-toxic at the concentration discharged into the receiving waters; however, the colour they impart is very undesirable to the water user for aesthetic reasons. Some dyes discharged into rivers are harmful to the aquatic life (Denizli et al., 2000; Acosta et al., 2005) because some of these effluent streams contain mixtures of heavy metal ions (i.e. Cr, Al, Cu,…) and other toxic organic pollutants including poly vinyl alcohol, starches, various surfactants, pesticides, biocides and others, all of which have varying Biochemical Oxygen Demand (BOD) (McKay et al.,1986; Werner et al., 2005). Also, the resulting wastewaters are highly toxic as some of them contain a large amount of metal complexing dyes (Denizli et al., 2000). Dyes often contain elements such as nitrogen, chlorine or sulphur. The oxidation products of these molecules may be more toxic than the parent molecule. Many of the dyes are, however, difficult to remove, as they are stable to light and heat and are biologically non-degradable.

Most of the methods that have been studied have focused on producing clean water suitable for recycling. There are many possible methods of degrading dyes, including adsorption (Malmary et al.,1985; Yeh et al., 1995; Allen et al., 2004), ozonation (Kawakami et al., 1978; Muthukumar et al., 2004), Fenton's oxidation (Rathi et al., 2003), electrochemical treatment (Lin et al., 1994; Vlysside et al., 2000), coagulation/flocculation (Do et al., 1994), biological treatment (Fu et al., 2002) and photodegradation (Davis et al., 1994; Tang et al., 1995). Colour removal from textile wastewaters, however, is difficult using a single conventional chemical or biological method (Kurbus et al., 2003). Users of dyes employ various methods to reduce the impact of waste arising from the dyeing process.

This paper describes the photodegradation of the dyes (Acid Blue 29 and Ethyl Violet in the presence/absence of NaOH and aluminium ion). Model solutions of dyes were used in the study to mimic the industrial dye effluent. The aim of the work was to optimise the photolytic and photocatalytic degradation of dyes in the presence of sodium hydroxide and aluminium ions.

2. Experimental

2.1. Apparatus and materials

A reaction vessel (Figure 1) of 100 ml capacity (A) was placed within the UV-lamp coil (40W, Philps) (B). The

*Corresponding author: chungchih.chen@itri.org.tw

whole system was enclosed in an aluminium box (C) to minimise the loss of UV radiation. TiO_2 as a catalyst and hydrogen peroxide as an oxidant was supplied by BDH. Sodium hydroxide was supplied by BDH. Acid Blue 29 and Ethyl Violet were of laboratory reagent grade (BDH) which were used without further purification. Distilled water was also used throughout the work.



Figure 1 Photodegradation reaction vessel.

2.2 Photolysis experiments

Four different experimental conditions: 1) aqueous dye solution, 2) dye solution containing Al^{3+}/NaOH, 3) dye solution containing TiO_2, and 4) dye solution containing H_2O_2, were investigated. A solution of 100 ppm of the dye was prepared using distilled water. In the case of a solution containing NaOH and Al^{3+}, 100 ml of the distilled water was replaced by 100 ml 2 M NaOH solution containing 60 ppm of standard aluminium solution ($Al((NO)_3)_3$, BDH). The reaction mixture was then stirred. Two sets of experiments were carried out to investigate the effect of TiO_2 as a catalyst and 30% H_2O_2 as an oxidant, on the degradation of dye solutions. TiO_2 as a catalyst was added to the dye solution in three different quantities, 0.1 g, 0.2 g, and 0.3 g and 5 ml. In a second set of experiment, 10 ml, and 15 ml of 30% H_2O_2, were added as an oxidant. The vessel was placed into the UV-lamp coil. In the case of TiO_2, the samples were centrifuged to remove the catalyst before the analyses. Samples were taken at hourly intervals for six hours. Degradation of a solution containing 50 ppm Acid Blue 29 and 50 ppm of Ethyl Violet in 2 M NaOH was also investigated using a quantity of 0.1 g TiO_2, 5 ml H_2O_2 and 60 ppm of standard aluminium solution respectively. All the experiments were carried out in duplicate. The degradation was monitored by UV/vis spectroscopy (Perkin Elmer, Model No. Lambda 9). The percentage of dye degradation was calculated from the UV spectroscopy peak height data.

3. Results and Discussion

3.1 Photodegradation of Acid Blue 29

3.1.1 Effect of Al and NaOH

From Table 1, it can be seen that the presence of aluminum and sodium hydroxide can dramatically increase the degradation of Acid Blue 29. After photolysis for one hour, 97% degradation of the Acid Blue 29 is achieved in the presence of aluminium and sodium hydroxide, but only 58% of the dye degraded in the absence of aluminium and sodium hydroxide. After four hours, 98% degradation of the dye is achieved in the absence of aluminium and sodium hydroxide. This shows that photolysis of the NaOH solution provides sufficient hydroxyl radicals to destroy Acid Blue 29. Kurbus reported that NaOH has a positive effect in a suitable concentration and can accelerate the degradation of the dye (Kurbus et al., 2003). Additionally, at sufficiently high pH values the formation of oxidizing species such as the oxide radical anion $O^{-\bullet}$, which reacts in a different manner from $OH^{\bullet}$,

can also be responsible for the rate enhancement of degradation (O'Shea et al., 1995). Although the dye solution containing aluminium ion which is present as $Al(OH)_4^-$ at high pH can retard the photodegradation (Grimes et al., 2000).

Table 1 Effect of Al and NaOH on degradation of Acid Blue 29

Time (h)	Degradation of Acid Blue 29 (%)	
	Absence of Al+NaOH	Presence of Al+NaOH
1	58	97
2	85	100
3	95	100
4	98	100
5	98	100
6	98	100

3.1.2 Effect of TiO_2

The results in Table 2 reveal that the addition of TiO_2 increases the degradation of Acid Blue 29; however, increasing the amount of TiO_2 reduces the degradation of the dye. In the presence of 0.1g of TiO_2, 96% degradation of the dye was achieved in two hours whereas in the absence of TiO_2 only 85% of the dye was degraded. The addition of 0.2 g and 0.3 g of TiO_2, however, degraded 93% and 85% of the dye respectively. The dye is virtually destroyed in the first hour in the presence of 0.1g, 0.2g and 0.3g of TiO_2 (Table 2). The degradation of 97% is also achieved in the absence of TiO_2 in the first hour. Sodium hydroxide provides $OH^{\bullet}$ radicals to increase the degradation of the dye (Majcen et al., 1997) and hence, complete degradation is achieved which shows that the presence of NaOH has significant effect.

Table 2 Effect of TiO_2 on degradation of Acid Blue 29.

Time (h)	Degradation of Acid Blue 29 (%)							
	Absence of NaOH+Al				Presence of NaOH+Al			
	Absence of TiO_2	Presence of TiO_2 (g)			Absence of TiO_2	Presence of TiO_2 (g)		
		0.1	0.2	0.3		0.1	0.2	0.3
1	58	79	77	77	97	100	100	100
2	85	96	93	85	100	100	100	100
3	95	98	97	97	100	100	100	100
4	98	-	-	-	100	-	-	-
5	98	-	-	-	100	-	-	-
6	98	-	-	-	100	-	-	-

3.1.3 Effect of H_2O_2

Increasing the concentration of H_2O_2 decreases the degradation as shown in Table 3. In the absence of H_2O_2, degradation of 98% is achieved after four hours, while the corresponding values in the presence of H_2O_2 are 81% (5 ml), 61% (10 ml) and 52% (15 ml). This shows that the addition of H_2O_2 retards the degradation of dye (Majcen et al., 1995). In the absence of H_2O_2 there is 97% degradation in one hour, but with the addition of 5, 10 and 15 ml the degradation is 70%, 45% and 70% respectively after one hour (Table 3). This shows that although H_2O_2 can affect the degradation, NaOH can still play an important role in assisting dye destruction. Kurbus and coworkers reported that the presence of H_2O_2 and NaOH increases the photodegradation but the presence of excess NaOH decreases the speed of the reaction (Kurbs et al., 2003). Higher concentrations of peroxide and NaOH can reduce the degradation of dyes because Na_2O_2 is produced and formation of $OH^{\bullet}$ is decreased as shown in equation 1 (Kurbus et al., 2002).

$$2NaOH + H_2O_2 + 6H_2O \rightarrow Na_2O_2 + 8H_2O \quad (1)$$

Table 3 Effect of H_2O_2 on degradation of Acid Blue 29

Time (h)	Degradation of Acid Blue 29 (%)							
	Absence of NaOH+Al				Presence of NaOH+Al			
	Absence of H_2O_2	Presence of H_2O_2 (ml)			Absence of H_2O_2	Presence of H_2O_2 (ml)		
		5	10	15		5	10	15
1	58	32	28	25	97	70	45	70
2	85	53	42	36	100	88	76	85
3	95	69	53	45	100	93	91	94
4	98	81	61	52	100	96	95	98
5	98	89	69	61	100	97	95	100
6	98	94	74	68	100	97	97	100

3.2 Photodegradation of Ethyl Violet

3.2.1 Effect of Al and NaOH

The data in Table 4 show that the presence of Al and NaOH increases the degradation of Ethyl Violet. On the other hand, in the absence of Al and NaOH, only 91% degradation is achieved after six hours which indicates that NaOH has positive effect on the rate of reaction in the presence of aluminium ion.

Table 4 Effect of Al and NaOH on degradation of Ethyl Violet

Time (h)	Degradation of Ethyl Violet (%)	
	Absence of Al+NaOH	Presence of Al+NaOH
1	11	98
2	11	-
3	32	-
4	85	-
5	91	-
6	91	-

3.2.2 Effect of TiO_2

The results in Table 5 show that although the addition of TiO_2 increases the degradation, increasing the quantities of TiO_2 decreases the degradation rate. After four hours a degradation of 99%, 87% and 82% is achieved in the presence of 0.1, 0.2 and 0.3 g of TiO_2 respectively. The degradation is almost complete after six hours but the absence of TiO_2 decreases the degradation to 91%. In the presence of TiO_2, NaOH and Al, the degradation of the Ethyl Violet can be completed in the first hour (Table 5). Degradation of 98% is achieved in the absence of TiO_2 in the first hour, which suggests that TiO_2 is not necessary as a catalyst in the degradation because the addition of NaOH provides hydroxyl radicals.

Table 5 Effect of TiO_2 on degradation of Ethyl Violet

Time (h)	Degradation of Ethyl Violet (%)							
	Absence of NaOH+Al				Presence of NaOH+Al			
	Absence of TiO_2	Presence of TiO_2 (g)			Absence of TiO_2	Presence of TiO_2 (g)		
		0.1	0.2	0.3		0.1	0.2	0.3
1	11	32	28	11	98	100	100	100
2	11	45	30	27	-	-	-	-
3	32	90	85	80	-	-	-	-
4	85	99	87	82	-	-	-	-
5	91	99	97	96	-	-	-	-
6	91	99	99	99	-	-	-	-

3.2.3 Effect of H_2O_2

Increasing the quantity of H_2O_2 from 5 ml to 15 ml in the dye solution increases the degradation as shown in

Table 6. Almost complete degradation of the dye is achieved within three hours but in the absence of H_2O_2, only 91% degradation is achieved after five hours. The data indicate that H_2O_2 supplies $OH^{\bullet}$ radicals to degrade the Ethyl Violet. Complete degradation of dye is achieved within one hour (Table 6) but increasing the quantity of H_2O_2 reduces the degradation because of the higher concentrations of peroxide and NaOH leading to the formation of Na_2O_2 as shown in equation 1. This result proves that the H_2O_2 is not essential as an oxidant for the decomposition of the Ethyl Violet in the presence of NaOH and Al.

Table 6 Effect of H_2O_2 on degradation of Ethyl Violet

Time (h)	Degradation of Ethyl Violet (%)							
	Absence of NaOH+Al				Presence of NaOH+Al			
	Absence of H_2O_2	Presence of H_2O_2 (ml)			Absence of H_2O_2	Presence of H_2O_2 (ml)		
		5	10	15		5	10	15
1	11	28	31	38	98	100	98	97
2	11	60	86	90	-	-	-	-
3	32	94	97	100	-	-	-	-
4	85	99	100	100	-	-	-	-
5	91	99	100	100	-	-	-	-
6	91	99	100	100	-	-	-	-

3.3 Photodegradation of a mixture of Acid Blue 29 and Ethyl Violet

3.3.1 Effect of Al and NaOH

The results in Table 7 indicate that the majority of the dye can be destroyed in one hour in the presence 2 M NaOH. Almost complete decomposition of the dye is achieved in six hours in the absence of NaOH and Al. When 60 ppm of the standard aluminium solution ($Al(NO_3)_3$) is added to the dye solution; however, the degradation is retarded. Only 79% degradation of the dyes is achieved in six hours of reaction time. Thus, these results confirm that it is the ease of free hydroxyl radical formation in the presence of NaOH which ultimately increases the degradation rate. In the presence of aluminium ion however the degradation of dye is retarded (Grimes et al., 2000).

Table 7 Effect of Al and NaOH on degradation of mixed dyes

Time (h)	Degradation of a mixture of 50ppm Acid Blue 29 and 50ppm Ethyl Violet (%)		
	Blank	2M NaOH	60ppm Al
1	31	96	18
2	58	97	30
3	78	100	39
4	89	-	55
5	95	-	66
6	98	-	79

3.3.2 Effect of TiO_2

In the presence of TiO_2, 99% degradation is achieved in three hours (Table 8). In the presence of NaOH and TiO_2, and NaOH, Al and TiO_2 99% degradation is achieved in the first hour. Whereas, in the presence of Al and TiO_2 100% degradation is achieved in five hours. This result reveals that the presence of NaOH increases the degradation of dye because it provides sufficient hydroxyl radicals to decompose the dyes.

Table 8 Effect of TiO_2 on degradation of mixed dyes

Time (h)	Degradation of a mixture of 50ppm Acid Blue 29 and 50ppm Ethyl Violet (%)			
	TiO_2	$NaOH+TiO_2$	$Al+TiO_2$	$NaOH+Al+TiO_2$
1	70	99	65	99
2	88	99	56	-
3	99	99	40	-
4	-	-	49	-
5	-	-	100	-
6	-	-	100	-

3.3.3 Effect of H_2O_2

Degradation of 86% of the mixture of dyes is achieved in one hour in the presence of $NaOH+Al+H_2O_2$ as compared to 50% (H_2O_2), 48% ($NaOH+H_2O_2$) and 41% ($Al+H_2O_2$). However the dyes are almost completely degraded in three hours in the presence of $NaOH+Al+H_2O_2$ as compared to five hours with $NaOH+ H_2O_2$. As mentioned earlier sodium hydroxide increased the rate of degradation, however the presence of excess H_2O_2 decreases the speed (equation 3). The degradation rate is in the order of $NaOH+Al+H_2O_2 > NaOH+H_2O_2 > Al+ H_2O_2 > H_2O_2$. There is decrease in the rate of degradation when the photolysis is carried out in the presence of Al^{3+} and H_2O_2 compared to when NaOH and H_2O_2 are used.

Table 9 Effect of H_2O_2 on degradation of mixed dyes

Time (h)	Degradation of a mixture of 50ppm Acid Blue 29 and 50ppm Ethyl Violet (%)			
	H_2O_2	$NaOH+H_2O_2$	$Al+H_2O_2$	$NaOH+Al+H_2O_2$
1	50	48	41	86
2	63	78	59	94
3	74	91	72	97
4	82	95	81	98
5	88	97	88	98
6	92	97	94	98

4. Conclusion

The results obtained using the photodegradation system show that Acid Blue 29 and Ethyl Violet can be degraded in the presence of sodium hydroxide. The addition of titanium dioxide as a catalyst and hydrogen peroxide as an oxidant in this system to enhance the degradation rate is not necessary. The developed system was also successfully used for the degradation of dyes in the presence of aluminium ions. These results show that the developed method can be applied to an industrial sample without the need for any catalyst or oxidant.

References

Acosta, D.R., Martinez, A.I., Lopez, A.A., and Magana, C.R. (2005). Titanium dioxide thin films: The effect of the preparation method in their photocatalytic properties. Journal of Molecular Catalysis A: Chemical. 228, 273-283.

Allen, S.J., McKay, G. and Porter, J.F. (2004). Adsorption isotherm models for basic dye adsorption by peat in single and binary component systems. Journal of Colloid and Interface Science. 280, 322-333.

Davis, R.J., Gainer, J.L., ONell, G. and Wu, I.W. (1994). Photocatalytic decolorization of wastewater dyes. Water Environment Research. 66, 50-53.

Denizli, A., Say, R. and Arica, Y. (2000). Removal of heavy metal ions from aquatic solutions by membrane chromatography.Seperation and Purification Technology. 21, 181-190.

Do, J.S. and Chen, M.L. (1994). Decolourization of dye-containing solutions by electrocoagulation. Journal of Applied Electrochemistry. 24, 785-790.

Fu, Y. and Viraraghavan, T. (2002). Dye biosorption sites in Aspergillus niger. Bioresource Technology. 82, 139-145.

Grimes, S.M., Donaldson, J.D., Chaudhary, A.J. and Hassan M. (2000) Simultaneous recovery of metals and destruction of organic species: Cobalt and phthalic acid. Environmental Science and Technology. 34, 4128-4132.

Kawakami, W., Hashimoto, S., Nishimura, K., Miyata, T. and Suzuki, N. (1978). Electron-beam oxidation treatment of a commercial dye by use of a dual-tube bubbling column reactor. Environmental Science and Technology. 12, 189.

Kurbus, T., Slokar, Y.M and Marechal, A.M. (2002). The study of the effects of the variables on H_2O_2/UV decoloration of vinylsulphone dye: Part Ⅱ. Dyes and Pigments. 54, 67-78.

Kurbus, T., Slokar, Y.M., Marechal, A.M. and Vonina, D.B. (2003). The use of experimental design for the evaluation of the influence of variables on the H_2O_2/UV treatment of model textile waste water. Dyes and Pigments. 58, 171-178.

Lin, S.H. and Peng, C.F. (1994). Treatment of textile wastewater by electrochemical method. Water Research. 28(2), 277-282.

Majcen, L.M., Slokar, Y.M. and Tanger, T. (1997). Decoloration of chlorotriazine reactive azo dyes with H_2O_2/UV. Dyes and Pigments. 33, 281-298.

Malmary, G., Perineau, F., Molinier, J. and Gaset, A. (1985). Continuous process of dye removal from liquid effluents using carbonized wool waste. Journal of Chemical Technology and Biotechnology Chemical Technology. 35A, 431-437.

McKay, G., Ramprasad, G. and Pratapa Mowli, P. (1986). Equilibrium studies for the adsorption of dyestuff from aqueous solutions by low-cost materials. Water Air and Soil Pollution. 29, 273-283.

Muthukumar, M. and Selvakumar, N. (2004). Studies on the effect of inorganic salts on decolouration of acid dye effluents by ozonation. Dyes and Pigments. 62, 221-228.

O'Shea K.E. and Cardona, C. (1995). The reactivity of phenol in irradiated aqueous suspensions of TiO_2. Mechanistic changes as a function of solution pH. Journal of Photochemistry and Photobiology A: Chemistry. 91, 67-72.

Rathi, A., Rajor, H.K. and Sharma, R.K. (2003). Photodegradation of direct yellow-12 using UV/H_2O_2/Fe^{2+}. Journal of Hazardous Materials. B102, 231-241.

Tang, W.Z. and An, H. (1995). UV/TiO_2 photocatalytic oxidation of commercial dyes in aqueous solutions. Chemosphere. 31(9), 4157-4170.

Vlyssides, A.G., Papaioanou, D., Loizidoy, M., Karlis, P.K. and Zorpas, A.A. (2000). Testing and Electrochemical method for treatment of textile dye wastewater. Waste Management. 20, 569-574.

Werner, J.J., McNeill, K. and Arnold, W.A. (2005). Environmental photodegradation of mefenamic acid. Chemosphere. 58, 1339-1346.

Yeh, R. and Thomas, A. (1995). Color difference measurement and color removal from dye wastewaters using different adsorbents. Journal of Chemical Technology and Biotechnology Chemical Technology. 63, 55-99.